

Aspiration pneumonia

Microbiology

- Aspiration pneumonitis is a sterile chemical inflammatory process caused by aspiration of gastric acid. Antibiotics are NOT indicated in aspiration pneumonitis.
- The vast majority of aspiration pneumonia cases are indistinguishable from CAP and HAP – with respect to presentation, microbiology, and therapy.
- Classic aspiration pneumonia has a protracted onset with putrid sputum and often a lung abscess or empyema. Anaerobes play a larger role in classic aspiration pneumonia.

Diagnosis

- The diagnosis of pneumonia is based on suggestive clinical features (cough, fever, sputum production, pleuritic chest pain, dyspnea) **AND** a new chest x-ray infiltrate.
- Pneumonia takes several *days* to develop after an aspiration event.
- All patients with suspected aspiration pneumonia should have a chest X-ray. Sputum culture should be collected if risk factors for antibiotic resistant organisms are present.

EMPIRIC TREATMENT

Classification		Duration (days)
Witnessed aspiration event	NONE – monitor patient If therapy initiated and patient improves rapidly (within 24-48 hours), aspiration pneumonitis is likely. Empiric antibiotics can be discontinued. If symptoms persist >48 hrs then consider treatment for aspiration pneumonia.	0
Aspiration pneumonitis	NONE – monitor patient If symptoms persist >48 hrs then consider treatment for aspiration pneumonia	0
Aspiration pneumonia <i>Anaerobes unlikely</i> • Majority of cases lacking criteria below	ceftriaxone 1 g IV q24h <i>If severe illness:</i> piperacillin-tazobactam 3.375 g IV q6h ADD vancomycin ¹ if known or suspected MRSA <i>If severe beta-lactam allergy:</i> moxifloxacin 400 mg IV/PO q24h	5-7
Classic aspiration pneumonia <i>Anaerobes likely:</i> • subacute onset • putrid sputum • lung abscess, necrotizing pneumonia, or empyema	ceftriaxone 1 g IV q24h PLUS metronidazole 500 mg PO/IV q12h <i>If severe illness:</i> piperacillin-tazobactam 3.375 g IV q6h ADD vancomycin ¹ if known or suspected MRSA <i>If severe beta-lactam allergy:</i> moxifloxacin 400 mg IV/PO q24h	≥7 Call Resp and/or ID

For patients with renal insufficiency, see “Antimicrobial dosing in renal insufficiency”

¹. For vancomycin dosing, refer “Vancomycin Dosing and Therapeutic Monitoring” chapter

Oral Step-Down and Duration

- Potential step-down regimen: amoxicillin-clavulanate 875/125 mg PO BID
If severe beta-lactam allergy: moxifloxacin 400 mg PO daily
- Classic anaerobic aspiration pneumonia involves lung abscess, necrotizing pneumonia, or empyema, and requires prolonged therapy. Respiriology and/or Infectious Diseases consultation recommended.